

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

---

Papers in Microbiology

Papers in the Biological Sciences

---

January 1974

## Sporulation of *Bacillus thuringiensis* Without Concurrent Derepression of the Tricarboxylic Acid Cycle

Kenneth W. Nickerson

University of Nebraska-Lincoln, [knickerson1@unl.edu](mailto:knickerson1@unl.edu)

John De Pinto

Northern Regional Research Laboratory, Agricultural Research Service, U.S. Department of Agriculture, Peoria, Illinois

Lee A. Bulla Jr.

Northern Regional Research Laboratory, Agricultural Research Service, U.S. Department of Agriculture, Peoria, Illinois

Follow this and additional works at: <https://digitalcommons.unl.edu/bioscimicro>



Part of the [Microbiology Commons](#)

---

Nickerson, Kenneth W.; De Pinto, John; and Bulla, Lee A. Jr., "Sporulation of *Bacillus thuringiensis* Without Concurrent Derepression of the Tricarboxylic Acid Cycle" (1974). *Papers in Microbiology*. 55.

<https://digitalcommons.unl.edu/bioscimicro/55>

This Article is brought to you for free and open access by the Papers in the Biological Sciences at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Papers in Microbiology by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

## Sporulation of *Bacillus thuringiensis* Without Concurrent Derepression of the Tricarboxylic Acid Cycle

KENNETH W. NICKERSON,<sup>1</sup> JOHN DE PINTO,<sup>2</sup> AND LEE A. BULLA, JR.<sup>3</sup>

Northern Regional Research Laboratory, Agricultural Research Service, U.S. Department of Agriculture, Peoria, Illinois 61604

Received for publication 17 September 1973

*Bacillus thuringiensis* sporulates in a glucose-glutamate medium without concurrent derepression of the tricarboxylic acid cycle. Glutamate appears to regulate tricarboxylic acid cycle activity as well as to influence spore heat resistance and production of dipicolinic acid.

Bacterial sporulation is a model system for studying cellular differentiation. However, biochemical and genetic characterization of spore formation has been plagued by difficulty in distinguishing between causal necessity and fortuitous coincidence of physiological events (7). Some events implicated as requisite for sporulation, because of their temporal relationship to morphological development or because of their absence in certain mutants unable to sporulate, may not be involved in spore formation at all. One such physiological phenomenon in aerobic sporeformers is the appearance of tricarboxylic acid cycle activity during post-logarithmic growth and sporogenesis.

Yousten and Hanson (13) provided the first evidence that the tricarboxylic acid cycle may not be an absolute requirement for sporulation. They showed that mutants of *Bacillus subtilis* with lesions in the tricarboxylic acid cycle could be made to sporulate by appropriate manipulation of the culture media. In this communication, we provide direct evidence that a fully operational tricarboxylic acid cycle is not necessarily required for sporulation.

We developed a chemically defined growth medium in which *B. thuringiensis* var. *entomocidus* NRRL B-4046 has sporulated repeatedly without an active tricarboxylic acid cycle. This medium consists of (per liter of distilled water): glucose, 1.0 g;  $(\text{NH}_4)_2\text{SO}_4$ , 2.0 g;  $\text{K}_2\text{HPO}_4$ , 0.5 g;  $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ , 0.3 g;  $\text{MnSO}_4 \cdot \text{H}_2\text{O}$ , 0.05 g;  $\text{CaCl}_2 \cdot 2\text{H}_2\text{O}$ , 0.08 g;  $\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$ , 0.005 g;  $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$ , 0.005 g;  $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$ , 0.0005 g; and glutamate, 0.16 g

(pH adjusted to 7.3 with KOH). The glutamate concentration is critical. This medium contains 0.016% glutamate and is designated LG (low glutamate), in contrast to our high glutamate (HG) medium which contains 0.2% glutamate. No growth occurs in the absence of glutamate. Refractile spores were formed in both the LG medium (Fig. 1) and the HG medium, but whereas the final pH of the HG medium was above 8.0, the pH of the LG medium dropped to about 4.5 and remained there throughout sporulation.

A final pH of 4.5 in the LG medium is presumptive evidence that the tricarboxylic acid cycle was not activated. This conclusion was confirmed by comparative radiorespirometric studies of sporulating cells from HG and LG media. Figure 2 shows the pattern of  $^{14}\text{CO}_2$  released from specifically labeled glutamate and acetate. Sporulating cells grown in the HG medium actively metabolized glutamate;  $^{14}\text{CO}_2$  release in the order  $\text{C2} = \text{C5} > \text{C3,4}$  is consistent with glutamate oxidation via an intact tricarboxylic acid cycle. In contrast, sporulating cells grown in LG medium did not oxidize either labeled glutamate or labeled acetate. Virtually no  $^{14}\text{CO}_2$  was released from either substrate. These results provide strong evidence that a functional tricarboxylic acid cycle was absent during sporulation in the LG medium because: (i) glutamate was incorporated into the sporulating cells during radiorespirometry; e.g., 20% of the glutamic acid-3,4- $^{14}\text{C}$  was incorporated into the cells, whereas only 0.23% was oxidized to  $\text{CO}_2$ ; (ii) separate uptake experiments showed that the concentration of carrier glutamate used in the radiorespirometry (0.02% in all experiments) was sufficient to ensure entry of the labeled glutamate into the cells; and (iii) by identical techniques, an active tricarboxylic acid cycle was demon-

<sup>1</sup> Present address: Department of Plant Pathology, University of Nebraska, Lincoln, Neb. 68503.

<sup>2</sup> On sabbatical leave from the Department of Biology, Bradley University, Peoria, Illinois 61606.

<sup>3</sup> Present address: U.S. Grain Marketing Research Center, U.S. Department of Agriculture, Manhattan, Kan. 66502.

strated in sporulating cells grown in HG medium.

Table 1 compares the properties of spores produced in HG and LG media. The LG-grown spores formed in the absence of a functional tricarboxylic acid cycle were: (i) refractile; (ii) frequently of normal *B. thuringiensis* spore morphology (see Fig. 1); (iii) octanol resistant (6); (iv) resistant to 62 C for 1 h; and (v) not stained by fat (4) and metachromatic granule (2) specific stains. The spores differed from those produced in HG media because they were: (i) sensitive to heat shock (30 min) at 80 C; (ii)

reduced in dipicolinic acid (8) content; and (iii) of decreased density. These properties are similar to those of the dipicolinic acid-deficient spores produced by dipicolinic acidless mutants (12) and by cells grown in the presence of ethyl oxamate (6), picolinamide (11), nicotinamide (9), or excess phenylalanine (5). Our work indicates that glutamate, either directly or indirectly, regulates dipicolinic acid production and heat resistance. Lack of a functional tricarboxylic acid cycle may be important to




FIG. 1. Phase contrast micrograph of *B. thuringiensis* spores formed in low glutamate medium.




FIG. 2. Radiospirometric patterns of glutamate and acetate utilization by *B. thuringiensis* var. *entomocidus*. Sporulating cells grown in high and low glutamate media were assayed by the procedures of Bulla et al. (3). Glutamate-2-<sup>14</sup>C; -3,4-<sup>14</sup>C; and -5-<sup>14</sup>C (0.02% glutamate) at pH 7. Acetate-1-<sup>14</sup>C; and -2-<sup>14</sup>C (0.003% acetate) at pH 4.6.

TABLE 1. Characteristics of spores formed in media of differing glutamate concentrations

Gluta- mate concn	Properties								
	Final pH of sporu- lation medium	Total viable count (per ml)	Octanol-re- sistant (6) count (per ml)	Heat-stable spores (%)		Dipicolinic acid con- tent (μg/10 <sup>8</sup> spores) <sup>a</sup>	Spore density <sup>b</sup> (g/ml)	Tricarboxylic acid cycle during sporu- lation	Glucose utilized during growth (%)
				80 C for 0.5 h	62 C for 1 h				
High	~8	1.8 × 10 <sup>8</sup>	1.9 × 10 <sup>7</sup>	100	100	6.7	1.30	Present	95.1
Low	4.5-4.8	2.8 × 10 <sup>8</sup>	2.8 × 10 <sup>7</sup>	<0.01	90-95	≤3.0 <sup>c</sup>	1.27	Absent	92.3

<sup>a</sup> Spores defined as octanol resistant counts.  
<sup>b</sup> Determined by Renografin density gradient centrifugation (12).  
<sup>c</sup> Highest of five determinations. Other values 2 to 2.5 μg/10<sup>8</sup> spores.

dipicolinic acid synthesis because aspartate is a precursor of dipicolinic acid.

Our data establish that the tricarboxylic acid cycle is inoperative during sporulation of *B. thuringiensis* var. *entomocidus* in LG medium; the presence or absence of individual tricarboxylic acid cycle enzymes was not established. The phenomenon of sporulation at low pH occurred also in *B. thuringiensis* var. *alesti* NRRL B-4041, the only other organism we tested.

High amino acid concentrations may induce enzymes for the catabolic degradation of that amino acid, and glutamate decarboxylase (L-glutamate l-carboxy-lyase EC 4.1.1.15) activity in *B. thuringiensis* can be high (1). J. N. Aronson (personal communication) has suggested that *B. thuringiensis* possesses a modified tricarboxylic acid cycle in which alpha-ketoglutarate dehydrogenase is absent. Instead, alpha-ketoglutarate is converted to glutamate and then to succinate via gamma-aminobutyric acid. We feel that high concentrations of glutamate may be necessary to induce glutamate decarboxylase or another enzyme of the gamma-aminobutyric acid pathway. This theory explains the high concentrations of glutamate required for sporulation in a great number of different organisms (7). Our radiorespirometric data (see also ref. 3) would not distinguish such a modified tricarboxylic acid cycle from one containing alpha-ketoglutarate dehydrogenase. At present, we are conducting enzyme assays to test these ideas.

This paper was presented, in part, at the 73rd Annual Meeting of the American Society for Microbiology, 6-11 May 1973, Miami Beach, Fla.

We gratefully acknowledge the capable technical assistance of Gordon L. Adams.

K. W. N. is a recipient of a postdoctoral resident research

associateship established by the Agricultural Research Service, U.S. Department of Agriculture, in association with the National Academy of Sciences, National Research Council, 1971-1973.

#### LITERATURE CITED

1. Akers, E., and J. N. Aronson. 1971. Detection on polyacrylamide gels of L-glutamic acid decarboxylase activities from *Bacillus thuringiensis*. *Anal. Biochem.* **39**:535-538.
2. Albert, H. 1920. Diphtheria bacillus stains with a description of a new one. *Amer. J. Pub. Health* **10**:334-337.
3. Bulla, L. A., G. St. Julian, and R. A. Rhodes. 1971. Physiology of sporeforming bacteria associated with insects. III. Radiorespirometry of pyruvate, acetate, succinate, and glutamate oxidation. *Can. J. Microbiol.* **17**:1073-1079.
4. Burdon, K. L. 1946. Fatty material in bacteria and fungi revealed by staining dried, fixed slide preparations. *J. Bacteriol.* **52**:665-678.
5. Church, B. D., and H. Halvorson. 1959. Dependence of the heat resistance of bacterial endospores on their dipicolinic acid content. *Nature (London)* **183**:124-125.
6. Gollakota, K. G., and H. O. Halvorson. 1963. Biochemical changes occurring during sporulation of *Bacillus cereus* T. II. Effect of esters of organic acids on sporulation. *J. Bacteriol.* **85**:1386-1393.
7. Hanson, R. S., J. A. Peterson, and A. A. Yousten. 1970. Unique biochemical events in bacterial sporulation. *Annu. Rev. Microbiol.* **24**:53-90.
8. Janssen, F. W., A. J. Lund, and L. E. Anderson. 1958. Colorimetric assay for dipicolinic acid in bacterial spores. *Science* **127**:26-27.
9. Mohankumar, K. C., and K. G. Gollakota. 1970. The effects of nicotinamide on heat resistance in spores of *Bacillus cereus* T. *Experientia* **26**:1162-1163.
10. Szulmajster, J., and R. S. Hanson. 1965. Physiological control of sporulation in *Bacillus subtilis*, p. 162-173. In L. L. Campbell and H. O. Halvorson (ed.), *Spores III*. American Society for Microbiology, Ann Arbor, Mich.
11. Upreti, G. C., R. P. Singh, J. Verma, P. L. Bhatia, and K. G. Gollakota. 1969. The effect of some alpha picolinic acid derivatives on growth and sporulation of bacilli. *Biochem. Biophys. Res. Commun.* **35**:611-618.
12. Wise, J., A. Swanson, and H. O. Halvorson. 1967. Dipicolinic acid-less mutants of *Bacillus cereus*. *J. Bacteriol.* **94**:2075-2076.
13. Yousten, A. A., and R. S. Hanson. 1972. Sporulation of tricarboxylic acid cycle mutants of *Bacillus subtilis*. *J. Bacteriol.* **109**:886-894.